Journal of Mammalogy, 85(6):1253–1254, 2004

INDEX TO VOLUME 85

AUTHORS

Abe, H., 396
Abramsky, Z., 67
Adrados, C., 774
Alves-Costa, C. P., 478
Amano, M., 311
Amman, B. R., 162
Ammerman, L. K., 842
Anderson, C. R., Jr., 1207
Anderson-Sprecher, R. C., 414
Arrizabalaga, A., 524
Arroyo-Cabrales, J., 133
Asher, M., 788
Ashley, M. C., 611
Aubry, K. B., 640
Avila-Flores, R., 675
Baker, B. W., 1145
Baker, R. J., 126
Baldwin, R. A., 1068
Bamshad, M., 1124
Banack, S. A., 43
Barbay, J. M., 331
Barnes, B. M., 260
Barrett, R. H., 649
Barrionuevo, F. J., 120
Barry, R. P., 932
Bashaw, M. J., 428
Batzli, G. O., 216
Bazykin, G., 105
Beck, T. D. I., 414
Bennett, M. B., 296
Bennett, N. C., 1225
Benson, J. F., 983
Bercovitch, F. B., 428
Bergstrom, B. J., 1077
Bertolino, S., 254
Bickham, J. W., 338
Blumstein, D. T., 25
Bogan, M. A., 995
Bonato, V., 708
Bonvicino, C. R., 316
Boulanger, J., 618
Bowyer, R. T., 498, 932, 1039
Bozinovic, F., 260
Bradley, R. D., 162, 389, 1160, 1184, 1194
Brady, M. J., 552
Braggio, E., 316
Breck, S. W., 1145
Brock, R. E., 51
Broders, H. G., 273
Browne, R. A., 875
Burgos, M., 120
Burns, J. A., 379
Busch, C., 290
Buskirk, S. W., 640
Câmara, B. G. O., 834
Câmara, E. V. C., 834
Cameron, G. N., 196
Canepuccia, A. D., 1111
Cardenas, R., 268
Carmona, A., 268
Carranza, J., 761
Carroll, D. S., 389, 1184
Casanave, E. B., 1111
Castro, A. C. S., 834
Cerchio, S., 963
Cervantes, F. A., 268
Chamberlain, M. J., 983
Chávez-Ramírez, F., 1095
Chesser, R. K., 58
Chimimba, C. T., 1225
Chowdhary, B. P., 924
Christófaro, C., 478
Clark, W. R., 187
Cluff, H. D., 576
Coertze, R. J., 756
Cook, J. A., 591, 601, 1149
Cook, J. G., 714
Cook, R. C., 714
Cotera, M., 1095
Coulson, G., 1053
Coutant, A. E., 245
Cowan, D. P., 959
Cryan, P. M., 995
Currado, I., 254
da Fonseca, G. A. B., 478
Dannelly, H. K., 15
Darling, J., 963
Dávalos, L. M., 404
Davis, H., 633
de Freitas, T. R. O., 1201
De Luna, E., 866
de Oliveira, J. A., 148
D'Elía, G., 897
Decker, D. J., 1215
Degen, A. A., 1106
Delgado, J., 268
Demarais, S., 768
Demboski, J. R., 591, 1170
Derr, J. N., 924
DeYoung, R. W., 768
Dickman, A. J., 19
Dizon, A. E., 347
Doan-Crider, D. L., 140
Dobson, F. S., 58
Doonan, T. J., 1102
du P. Bothma, J., 756
Dumont, E. R., 8
Duncan, N. P., 470
Durish, N. D., 1160
Eccard, J. A., 440
Eddingsaas, A. A., 601
Edge, W. D., 87
Edwards, C. W., 389
Eilam, D., 748
Eiler, K. C., 43
Elangovan, V., 806
Eldridge, D. J., 1060
Elias, S. P., 743
Engelken, T. J., 768
Escorza-Treviño, S., 347
Estrada, E., 1095
Findlay, C. S., 273
Flannery, T. F., 825
Flaquer, C., 524
Flores-Martínez, J. J., 133
Foltz, D. W., 58
Frame, P. F., 505
Franklin, W. L., 72
Frantz, A. C., 959
Gabriele, C. M., 963
Gamarra de Fox, I., 698
Garden, J. G., 296
Gava, A., 1201
Gehring, T. M., 79
Gehrt, S. D., 483
Gemmell, N. J., 338
Gende, S. M., 633
Genoways, H. H., 852
Gilchrist, J. S., 491
Givens, H. L., 331
Glockner-Ferrari, D. A., 963
Gomes Facure, K., 708
Gonçalves, P. R., 148
Gonzales, R. A., 768
González, D., 866
Gorresen, P. M., 688
Grant, W. E., 453
Greenwood, R. J., 187
Grewal, S. K., 625
Guerrero, J. A., 866
Guthrie, H. D., 768
Haberl, W., 396
Hadley, G. L., 97
Hafner, D. J., 1170
Hafner, M. S., 1170
Halbert, N. D., 924
Halcomb, K. E., 1160
Hamilton, M. J., 1184
Harlow, H. J., 414
Harris, J. H., 517
Haughland, D. L., 225
Havelka, M. A., 940
Hayano, A., 311
Hayes, L. D., 781
Haynie, M. L., 1184
Helgen, K. M., 825
Hellgren, E. C., 140
Herman, L. M., 963
Herrera, L. G., 133
Hertel, F., 886
Hintz, R. J., 446
Honeycutt, R. L., 453
Hoofer, S. R., 321
Hoogland, J. L., 58
Houston, A. E., 1068
Hunter, M. L., Jr., 743
Ibáñez, C., 38
Illoldi-Rangel, P., 658
Im, S., 25
Irwin, M. T., 420
Iwasa, M. A., 396
Jacobsen, B. K., 601
Jacquot, J. J., 1009
Janeau, G., 774
Jansa, S. A., 404
Janse van Rensburg, L., 1225
Jean-Baptiste, N., 1124
Jiménez, R., 120
Johnson, D. S., 932
Johnson, W. E., 432
Jorgensen, E. E., 531
Jurasz, C., 963
Juste, J., 38
Kalko, K. V., 1116
Kam, M., 1106
Kaufman, D. W., 562
Kaufman, G. A., 562
Kawamichi, T., 30
Keeley, A. T., 113
Keeley, B. W., 113
Kelt, D. A., 51
Kenagy, G. J., 260
Kennedy, M. L., 1068
Kester, C. L., 995

Khokhlova, I. S., 1106
Kie, J. G., 498
Kilpatrick, C. W., 389, 1160, 1184
Klein, D. R., 591
Kolosova, I., 723
Kruchek, B. L., 569
Kung, T. K., 625
Kyle, C. J., 633
Lambertsen, R. H., 446
Landis, G. P., 995
Larsen, K. W., 225
LaVal, R. K., 237
Lehman, N., 1215
Leitner, P., 517
Leopold, B. D., 983
Lessa, E. P., 601
Libhaber, N., 748
Light, J. E., 1170
Lin, Y. K., 216, 781
Linders, M. J., 511
Lindzey, F. G., 1207
Liu, P. S., 1068
Lohuis, T., 414
López-González, C., 989
López-Wilchis, R., 38
Loughlin, T. R., 338
Lucherini, M., 1111
Lucid, M. K., 1149
Luis, J., 268
Macdonald, D. W., 432
Maher, C. R., 1087
Main, M. B., 973
Maldonado, J. E., 886
Mandelli, M.-J., 282
Manfredi, C., 1111
Mann, C., 774
Manning, J. A., 87
Marimuthu, G., 1, 806
Marin, J. C., 384
Marker, L. L., 19
Martínez, M., 268
Mason, M. J., 797
Masuda, R., 396
Mattson, D. J., 731
May, B. P., 133
Mazzoglio, P. J., 254
McDonald, D. B., 1207
McLoughlin, P. D., 576
McPhee, M. E., 1130
Mech, L. D., 505, 714
Medellín, R. A., 196, 675
Meijaard, E., 302
Meikle, D. B., 1015
Meyer, J., 440
Michener, G. R., 1019
Millar, J. S., 940
Mizroch, S. A., 963
Moreno-Valdez, A., 453
Moshkin, M., 723
Muller, L. I., 768
Mullin, S. K., 911
Muñiz M., R., 1184
Munro, R., 618
Nelson, M. E., 505
Nesterenko, V. A., 396
Newhouse, N. J., 633
Nichols, J. V., 540, 663
Nicodemus, A., 25
Nishibayashi, K., 30
Nogueira, J. C., 834
Núñez-Garduño, A., 38
Nusser, S. M., 187
O'Brien, S. J., 432
Ohdachi, S. D., 396
Oi, T., 311
Olesen, C. R., 245
O'Neal, R., 8
Onorato, D. P., 140
Otali, E., 491
Palma, R. E., 384
Paquet, P. C., 576
Pardiñas, U. F. J., 897
Pastene, L. A., 347
Patton, J. C., 338
Penny, C. G., 428
Pépin, D., 774
Phillips, M. L., 187
Pillay, N., 911
Pocock, M. J. O., 959
Porter, C. A., 126
Prange, S., 483
Prentice, D. A., 15
Priya, S., 806
Pulido, F. J., 761
Raghuram, H., 806
Raharison, J.-L., 420
Ramp, D., 1053
Ramsay, M. A., 576
Randall, J. A., 723, 1002
Ransome, D. B., 206
Raudsepp, T., 924
Reeder, S. A., 1194
Rehmeier, R. L., 562
Rhymer, J. M., 331
Rieches, R. G., 428
Rodriguez, R. M., 842
Rogovin, K., 723
Rose, R. K., 1077
Rosell, F., 1138
Ruedi, M., 133
Russell, G. A., 640
Rye, R. O., 995
Sachser, N., 788
Salden, D. R., 963
Sales, G., 282
Samonds, K. E., 420
Sánchez-Cordero, V., 658
Sánchez-Prieto, C. B., 761
Sarno, R. J., 432
Schleich, C. E., 290
Schlexer, F. V., 649
Schmidt, G. A., 649
Schmidt, K. N., 649
Schnell, G. D., 948
Schoeman, A. S., 1225
Schulte, B. A., 1138
Scott-Morales, L., 1095
Searle, J. B., 959
Shami, K., 625
Shibata, F., 30
Shier, D. M., 1002
Singaravelan, N., 1
Sipe, T. W., 875
Slade, N. A., 552, 1102
Smith, W. P., 540, 663
Snider, I., 1106
Solomon, N. G., 781, 1009
Sorin, A. B., 356
Sovada, M. A., 187
Spinelli de Oliveira, E., 788
Spotorno, A. E., 384
Spradling, T. A., 1170
Stadelmann, B., 133
Stancampiano, A. J., 948
Stenhouse, G., 618
Stevens, R. D., 698
Stewart, B. S., 1215
Straley, J. M., 963
Strobeck, C., 633
Sugg, D. W., 58
Sullivan, T. P., 206
Sulok, M., 1102
Sundell, J., 440
Sunquist, M. E., 973
Swihart, R. K., 79
Taylor, P. J., 911
Tchabovsky, A., 105
Terleph, T. A., 1124
Theberge, J. B., 625
Theberge, M. T., 625
Theil, P. K., 245
Thornton, D. H., 973
Timm, R. M., 852
Torre, I., 524
Torres-Morales, L., 989
Townsend Peterson, A., 658
Truex, R. L., 649
Trujillo, R. G., 338
Tsurim, I., 67
Uieda, W., 708
Vaiana, M., 254
Valladares, J. P., 384
Van Den Bussche, R. A., 140, 321
van der Merwe, M., 1225
Vander Haegen, W. M., 511
Vázquez, L. B., 196
Vázquez, R. A., 260
Vilà, C., 886
von Ziegesar, O., 963
Waltari, E., 591
Walton, L. R., 576
Weinbeer, M., 1116
Weir, R. D., 633
Welch, G. R., 768
West, S. D., 511
Whitaker, J. O., Jr., 15, 460
White, B. N., 625
White, P. C. L., 959
Wiggers, E. P., 483
Wilder, S. M., 1015
Williams, C. L., 1145
Willig, M. R., 688, 698
Wilson, K. R., 97
Wilson, P. J., 625
Wisely, S. M., 640
Witham, J. W., 743
Wright, P. C., 420
Yamaguchi, N., 432
Young, J. K., 72
Yuvana, E., 806
Zheng, L., 273
Zielinski, W. J., 470, 640, 649
Zugmeyer, C., 25
Zuleta R., C., 384
Zurita, F., 120

Journal of Mammalogy, 85(6):1255–1264, 2004

SUBJECTS

(New names in *italic* type)

A
Abrocoma cinerea, 384
Abundance:
  Clethrionomys gapperi wrangeli, 540
  microhabitat affinities of small mammals, 948
  Peromyscus keeni macrorhinus, 540
  Peromyscus leucopus, 743
  Peromyscus maniculatus, 562
Acinonyx jubatus jubatus, 19
Acomys cahirinus, 1106
Acorns, as food source, 743
Activity patterns:
  Callosciurus finlaysonii, 254
  Capreolus capreolus, 245
  Octodon degus, 260
  Odocoileus hemionus, 498
  Thallomys nigricauda, 440
Adaptive radiation:
  Myotis vivesi, 133
Adaptive sex ratio theories, 768
Admiralty Islands:
  Spilocuscus kraemeri, 825
Africa:
  Acinonyx jubatus jubatus, 19
  Dasymys (4 spp.), 911
  Mungos mungo, 491
Age estimation:
  Cryptomys hottentotus pretoriae, 1225
Age structure:
  Cervus elaphus nelsoni, 714
  Sigmodon hispidus, 1077
Aggression, intraspecific:
  Lynx rufus, 983
  Microtus ochrogaster, 1124
Alarm call:
  Spermophilus (2 spp.), 43
Alaska:
  Clethrionomys gapperi wrangeli, 540
  Glaucomys sabrinus griseifrons, 663
  Peromyscus keeni, 1149
  Peromyscus keeni macrorhinus, 540
Alberta:
  Dicrostonyx groenlandicus, 379
  Lemmus trimucronatus, 379
Alexander Archipelago:
  Peromyscus keeni, 1149
Alien species:
  Callosciurus finlaysonii, 254
Allopatry:
  Equus caballus, 611
Allozymes:
  Cynomys ludovicianus, 58
American badger, 633, 1019, 1060
American marten, 470
American mink, 432
Amplified fragment length polymorphism (AFLP):
  Sorex (2 spp.), 875
Antarctic pinnipeds, 1215
Appalachian Mountains:
  Sorex (2 spp.), 331
Arboreality:
  Glirulus japonicus, 440
  Thallomys nigricauda, 440
Archonta, 321
Arctic, historical biogeography of, 591
Arctic ground squirrel, 601
Argentina:
  Ctenomys talarum, 290
  Oncifelis geoffroyi, 1111
Artibeus:
  fimbriatus, 688
  hirsutus, 866
  inopinatus, 866
  jamaicensis jamaicensis, 866
  jamaicensis paulus, 866
  jamaicensis richardsoni, 866
  jamaicensis triomylus, 866
  jamaicensis yucatanicus, 866
  lituratus, 688, 866
  obscurus, 866
  planirostris fallax, 866
Asia:
  Crocidura (8 spp.), 396
  Panthera pardus, 302
  Rhombomys opimus, 723
Asiatic black bear, 311
Atlantic rainforest:
  bat community ecology, 698
  bat habitat fragmentation, 688
Australia:
  Macropus giganteus, 1053
Aztec mouse, 196

B
Badger, 633, 1019, 1060
Baiomys:
  musculus, 162, 395, 658
  taylori, 162, 395
Balaenoptera acutorostrata, 446
Balantiopteryx plicata, 38, 658
Banded mongoose, 491
Bandicoot, northern brown, 296
Barn owl pellets, 524
Barro Colorado Island:
  Lampronycteris brachyotis, 1116
Bats:
  Artibeus (2 spp.), 688
  Artibeus jamaicensis, 658
  Balantiopteryx plicata, 38, 658
  Carollia perspicillata, 658, 688
  cave use by Mexican bats, 675
  Chrotopterus auritus, 688, 708
  climate change effects on, 237
  community ecology, 698
  Corynorhinus (2 spp.), 989
  Cynopterus sphinx, 1
  Desmodus rotundus, 658
  Dobsonia minor, 8
  Ectophylla alba, 126
  Eptesicus fuscus, 15 460
  Eumops (2 spp.), 852
  Glossophaga soricina, 658, 688
  habitat fragmentation, 688
  Lampronycteris brachyotis, 1116
  Lasionycteris noctivagans, 15
  Lasiurus (2 spp.), 15
  Lasiurus borealis, 460
  Lasiurus cinereus, 995
  Leptonycteris nivalis, 453
  Lochophylla (5 spp.), 404
  Mesophylla macconnelli, 126
  Myotis (2 spp.), 273, 842
  Myotis (3 spp.), 15, 460
  Myotis vivesi, 133
  Nycticeius humeralis, 15, 460
  Nyctimene albiventer, 8
  Paranyctimene raptor, 8
  Pipistrellus subflavus, 15, 460
  Pizonyx vivesi, 133
  Platyrrhinus lineatus, 688
  Pteropus conspicillatus, 8
  Pteropus giganteus, 1
  Pygoderma bilabiatum, 688
  Rousettus leschenaulti, 1, 806
  Sturnira lilium, 688
  Syconycteris australis, 8
  Tadarida braziliensis, 113
  Trachops cirrhosus, 708
  Vampyressa (4 spp.), 126
  Vampyriscus (2 spp.), 126
  Vampyrum spectrum, 708
Bears:
  Ursus americanus, 140, 414
  Ursus arctos, 618
  Ursus arctos horribilis, 731
  Ursus thibetanus japonicus, 311
Beavers:
  Castor canadensis, 1145
  Castor fiber, 1138
Behavior. *See also specific behaviors*
  burrow use in Dipodomys stephensi, 51
  Capreolus capreolus heart rate variation, 245
  latrine behavior in lemurs, 420
  sexual segregation, 1039
  social hibernation, 25
Behavioral isolation:
  Equus caballus, 611
Beringia:
  Lepus (3 spp.), 591
  Spermophilus parryii, 601
Biak-Supiori:
  Spilocuscus wilsoni, 825
Bias, tampering, 959
Big brown bat, 15, 460
Biogeography:
  Baiomys (2 spp.), 162
  Dicrostonyx groenlandicus, 379
  Lemmus trimucronatus, 379
  Mexican mammal distribution, 658
  of northern high latitudes, 591
  Panthera pardus, 302
  Sorex (3 spp.), 886
Biopedturbation, by badgers, 1060
Birth date, estimating, 932
Bismark Archipelago:
  Spilocuscus (2 spp.), 825
Bison bison, 924
Black bear, 140, 414
Black-tailed prairie dog, 58
Black-tailed tree rat, 440
Bobcat, 973, 983
Body condition:
  Cervus elaphus nelsoni, 714
  Mungos mungo, 491
Body fat:
  Cervus elaphus nelsoni, 714
Body mass:
  Cryptomys hottentotus pretoriae, 1225

Body size:
Dipodomys heermanni arenae, 1002
Sigmodon hispidus, 1077
Body temperature:
hibernating Ursus americanus, 414
Bonneted bat, 852
Brazil:
Cavia aperea, 788
Chironectes minimus, 834
Chrotopterus auritus, 708
Ctenomys minutus, 1201
Nasua nasua, 478
Trachops cirrhosus, 708
Vampyrum spectrum, 708
Brazilian free-tailed bat, 113
Breeding. *See also* Mating; Reproduction
Cynomys ludovicianus, 58
delayed in Meriones tamariscinus, 105
Tadarida braziliensis, 113
Brown-nosed coati, 478
Burrow-use behavior:
Dipodomys stephensi, 51

C
California:
Dipodomys heermanni arenae, 1002
Dipodomys stephensi, 51
Martes (2 spp.), 470
Martes pennanti, 649
Odocoileus hemionus, 498
Spermophilus mohavensis, 517
California myotis, 842
Callosciurus finlaysonii, 254
Canada:
Canis lupus, 576
Canis lycaon, 625
Dicrostonyx groenlandicus, 379
Glaucomys sabrinus, 206
Lemmus trimucronatus, 379
Peromyscus leucopus, 940
Sorex (2 spp.), 331
Spermophilus richardsonii, 1019
Tamiasciurus douglasii, 206
Tamiasciurus hudsonicus, 225
Taxidea taxus, 1019
Ursus arctos, 618
Canis:
latrans, 625, 973
lupus, 576, 625
lupus familiarus, 43
lycaon, 625
rufus, 625
Capreolus capreolus, 245
Captivity, effect of:
on Peromyscus polionotus subgriseus, 1130
Capture-recapture studies:
Megaptera novaeangliae, 963
minimum number alive (MNA) population indices, 959
Procyon lotor, 483
Capture success:
Mephitis mephitis, 1068
Carollia perspicillata, 688
Castor:
canadensis, 1145
fiber, 1138
Cave use by Mexican bats, 675
Cavia aperea, 788
Cervus:
elaphus, 774
elaphus hispanicus, 761
elaphus nelsoni, 714
Cetaceans:
Balaenoptera acutorostrata, 446
Megaptera novaeangliae, 963
Phocoenoides dalli, 347
Chaetodipus hispidus, 948
Cheetah, 19
Chile:
Lama guanicoe, 72
Octodon degus, 260
Chinchilla:
brevicaudata, 384
lanigera, 384
Chipmunk, least, 97
Chiroptera:
Artibeus (2 spp.), 688
Carollia perspicillata, 688
Chrotopterus auritus, 688
climate change effects on, 237
community ecology, 698
Corynorhinus (2 spp.), 989
Eptesicus fuscus, 460
Eumops floridanus, 852
Glossophaga sorcina, 688
habitat fragmentation, 688
Lasiurus borealis, 460
Lonchophyllini, 404
Myotis (2 spp.), 273
Myotis (3 spp.), 460
Myotis vivesi, 133
Nycticeius humeralis, 460
phylogenetics, 321
Pipistrellus subflavus, 460
Platyrrhinus lineatus, 688
Pygoderma bilabiatum, 688
Sturnira lilium, 688
Tadarida braziliensis, 113
Chitinase, in insectivorous bats, 15
Chrotopterus auritus, 688, 708
Clethrionomys:
gapperi, 97
gapperi wrangeli, 540
Climate change:
biogeography of northern high latitudes, 591
effect on tropical cloud forest bats, 237
Cline:
Sorex (2 spp.), 331
Cloud forest:
bat populations, 237
rodent diets, 196
Cluster analysis:
microhabitat affinities of small mammals, 948
Coarse data, 932
Coati, brown-nosed, 478
Colorado:
Clethrionomys gapperi, 97
Marmota flaviventris, 25
Microtus montanus, 97
Peromyscus maniculatus, 97
Phenacomys intermedius, 97
Sorex (2 spp.), 97
Spermophilus lateralis, 97
Tamias minimus, 97
Zapus princeps, 97
Communities:
bat habitat fragmentation, 688
grassland rodent community dynamics, 552
small mammal richness, 524
Community ecology:
Paraguay bats, 698
Conservation:
bat habitat fragmentation, 688
Bison bison, 924
Corynorhinus (2 spp.), 989
Cynomys mexicanus, 1095
Dipodomys stephensi, 51
Eumops floridanus, 852
sexual segregation, implications of, 1039
Taxidea taxus, 633
Control region genes:
Myotis (2 spp.), 842
Sorex (2 spp.), 875
Copulation strategies:
Tadarida braziliensis, 113
Corynorhinus:
mexicanus, 989
townsendii, 989
Costa Rica:
cloud forest bats, 237
Cotton rat, 552, 948, 1077, 1102
Cougar, 1207
Coyote, 625, 973
Crabeater seal, 1215
Craniometrics:
Artibeus (10 spp.), 866
Cryptomys hottentotus pretoriae, 1225
Dasymys (4 spp.), 911
Panthera pardus, 302
Cratogeomys:
fumosus, 1170
gymnurus, 1170
neglectus, 1170
planiceps, 1170
tylorhinus, 1170
zinseri, 1170
Creeping vole, 87
Crocidura:
dsinezumi, 396
gemelini, 396
kurodai, 396
lasiura, 396
shantungensis, 396
sibirica, 396
suaveolens, 396
watasei, 396
Cryptomys hottentotus pretoriae, 1225
Cryptotis parva, 948
Ctenomys:
minutus, 1201
sociabilis, 797
talarum, 290
Cuica d'água, 834
Cynomys:
ludovicianus, 58
mexicanus, 1095
Cynopterus sphinx, 1
Cytochrome-*b* gene sequences:
Artibeus (10 spp.), 866
Baiomys (2 spp.), 162
chinchilids, 384
Crocidura (8 spp.) shrews, 396
Lonchophyllini, 404
Myotis (2 spp.), 842
Myotis vivesi, 133
neotomine-peromycine rodents, 389
Oxymycterus (3 spp.), 148
Peromyscus (6 spp.), 1184
Peromyscus keeni, 1149
Peromyscus truei species group, 1160
Scapteromys (2 spp.), 897
Sorex (2 spp.), 875
Spermophilus parryii, 601
Thrichomys (4 spp.), 316
Vampyressa, 126

D
Daily travel distance:
Cervus elaphus, 774
Dall's porpoise, 347
Dall's sheep, 932
Dasymys:
foxi, 911

incomtus, 911
longipilosus, 911
rufulus, 911
Date of birth, estimating, 932
Deer:
Capreolus capreolus, 245
Cervus elaphus, 774
Cervus elaphus hispanicus, 761
Cervus elaphus nelsoni, 714
Odocoileus hemionus, 498
Odocoileus virginianus, 356, 505, 658, 768
Deer mouse, 87, 97, 552, 562, 948
Deer tick, 1015
Defecation:
latrine behavior in lemurs, 420
Degu, 260
Delayed implantation:
Mustela vison, 432
Demography:
Clethrionomys gapperi wrangeli, 540
Cynomys ludovicianus, 58
Megaptera novaeangliae, 963
Microtus ochrogaster, 781
Peromyscus keeni macrorhinus, 540
Tamiasciurus hudsonicus, 225
Denmark:
Capreolus capreolus, 245
Denning behavior:
Canis lupus, 576
Glaucomys sabrinus, 206
Tamiasciurus douglasii, 206
Dental anomalies:
Acinonyx jubatus jubatus, 19
Dentin matrix protein 1, 1194
Deuterium, 995
Developmental stages:
Talpa occidentalis, 120
Dialect, 43
Dicrostonyx groenlandicus, 379
Didelphidae:
Chironectes minimus, 834
Diet:
Martes (2 spp.), 470
Nasua nasua, 478
Oncifelis geoffroyi, 1111
Oryzomys palustris, 569
Peromyscus aztecus, 196
Reithrodontomys fulvescens, 196
Digestion:
chitinase in insectivorous bats, 15
Discriminant function analysis:
microhabitat affinities of small mammals, 948
Dispersal:
delayed in Meriones tamariscinus, 105
Microtus (2 spp.), 216
Microtus ochrogaster, 781, 1009
Tamiasciurus hudsonicus, 225
Distribution:
Cratogeomys gymnurus species group, 1170
Cynomys mexicanus, 1095
Dasymys (4 spp.), 911
Eumops floridanus, 852
modeling Mexican mammal, 658
Procyon lotor, 483
tropical cloud forest bats, 237
Divergence. *See also* Molecular divergence
Spermophilus parryii, 601
Diversity. *See also* Genetic diversity; Habitat diversity
bat, 98, 688, 698
D-loop sequences:
Sorex (2 spp.), 875
DNA genotyping:
Ursus arctos, 618
Dobsonia minor, 8
Domestic dog, 43
Douglas squirrel, 206
Downed wood, 87
DQα exon, 1215

E
Eastern grey kangaroo, 1053
Eastern pipistrelle, 15, 460
Eastern wolf, 625
Echolocation:
effect of clutter on call structure, 273
habitat variation and jamming avoidance in Balantiopteryx plicata, 38
Ecological constraints:
Microtus ochrogaster, 1009
Ecological niche modeling, 658
Ecological separation:
coyotes and bobcats, 973
Ecology:
Corynorhinus (2 spp.), 989
Ectophylla alba, 126
Elk, 714
Emigration:
Microtus (2 spp.), 216
Endangered species:
Dipodomys stephensi, 51
Eumops floridanus, 852
Endemic populations:
Clethrionomys gapperi wrangeli, 540
Peromyscus keeni, 1149
Peromyscus keeni macrorhinus, 540
Energy metabolism:
Capreolus capreolus, 245
England:
Mustela vison, 432
Eptesicus fuscus, 15, 460
Equus caballus, 611
Eumetopias jubatus, 338
Eumops:
floridanus, 852
glaucinus, 852
Eurasian beaver, 1138
European roe deer, 245
Evening bat, 15, 460
Evolutionary innovation:
Balaenoptera acutorostrata, 446

F
Fecal analysis:
Oncifelis geoffroyi, 1111
Feeding behavior:
food hoarding in gerbils, 67
Glaucomys sabrinus, 206
Myotis vivesi, 133
nectar-feeding by pteropodid bats, 1
old world fruit bats, 8
Tamiasciurus douglasii, 206
Field vole, 282
Finlayson's squirrel, 254
Fisher, 470, 640, 649
Fishing bat, 133
Flight performance:
Rousettus leschenaulti, 806
Florida:
Canis latrans, 973
Eumops floridanus, 852
Lynx rufus, 973
Flow cytometry, 768
Flying squirrel, 206, 663
Food availability:
Leptonycteris nivalis, 453
Peromyscus leucopus, 743
Food caches:
Thomomys talpoides, 731
Food habits:
Canis latrans, 973
Chrotopterus auritus, 708
Eptesicus fuscus, 460
Lasiurus borealis, 460
Lynx rufus, 973
Martes (2 spp.), 470
Myotis (3 spp.), 460
Nycticeius humeralis, 460
Oncifelis geoffroyi, 1111
Pipistrellus subflavus, 460
Trachops cirrhosus, 708
Ursus arctos horribilis, 731
Vampyrum spectrum, 708
Food hardness:
effect on fruit bat feeding, 8
Food hoarding:
in gerbils, 67
Food resources:
Procyon lotor, 483
Food supplementation:
Cervus elaphus hispanicus, 761
Glaucomys sabrinus, 206
Sigmodon hispidus, 1102
Tamiasciurus douglasii, 206
Foraging behavior:
Callosciurus finlaysonii, 254
Lampronycteris brachyotis, 1116
Martes (2 spp.), 470
Odocoileus hemionus, 498
Taxidea taxus, 1019, 1060
Fossil species:
Eumops floridanus, 852
Panthera pardus, 302
Fox, red, 187
Fractal dimension, 187
Fragmentation, of Cynomys mexicanus colonies, 1095
France:
Cervus elaphus, 774
Fulvous harvest mouse, 196
Functional diversity:
Paraguay bats, 698

G
Gait:
Isoodon macrourus, 296
Garbage dumps, 491
GARP (genetic algorithm for rule-set prediction), 658
Gaspé shrew, 331
Gender determination:
Castor canadensis, 1145
Gene flow:
Martes pennanti, 640
Puma concolor, 1207
Ursus americanus, 140
Genetic diversity:
Antarctic pinnipeds, 1215
Bison bison, 924
Canis lycaon, 625
Cynomys ludovicianus, 58
Martes pennanti, 640
Peromyscus keeni, 1149
Genetic structure:
Martes pennanti, 640
Puma concolor, 1207
Taxidea taxus, 633
Genetic variation:
Oxymycterus (3 spp.), 148
Scapteromys (2 spp.), 897
Sorex (3 spp.), 886
Genet scats, 524
Genital system:
Chironectes minimus, 834

Geoffroy's cat, 1111
Geographic distribution:
  Cratogeomys gymnurus species group, 1170
  Cynomys mexicanus, 1095
  Dasymys (4 spp.), 911
  modeling Mexican mammal, 658
Geographic Information Systems, 658
Geographic variation:
  alarm call variability, 43
  Eumetopias jubatus, 338
  Oncifelis geoffroyi, 1111
  Oxymycterus (3 spp.), 148
  Sorex (2 spp.), 331
  Ursus thibetanus japonicus, 311
Georgia:
  Sigmodon hispidus, 1077
Gerbillus:
  allenbyi, 67
  pyramidum, 67
Gerbils:
  food hoarding by, 67
  great, 723
  tamarisk, 105
Giraffa camelopardalis, 428
Giraffe, 428
Glaucomys:
  sabrinus, 206
  sabrinus griseifrons, 663
Glirulus japonicus, 30
Global warming:
  effect on tropical cloud forest bats, 237
Glossophaga sorcina, 658, 688
Golden-mantled ground squirrel, 43, 99
Goodnight, Charles, 924
GPS tracking:
  Cervus elaphus, 774
  Odocoileus virginianus, 505
Grasslands:
  predator movement, 187
  rodent community, 552
Grazing enclosures, 1053
Great gerbil, 723
Grizzly bear, 618, 731
Group size:
  Odocoileus hemionus, 498
  Rhombomys opimus, 723
Growth:
  Rousettus leschenaulti, 806
Guanacos, 72
Guinea pig, 788

H
Habitat boundaries:
  Microtus (2 spp.), 216
Habitat diversity:
  Dicrostonyx groenlandicus, 379
  Lemmus trimucronatus, 379
Habitat fragmentation:
  bat response to, 688
  Mustela frenata, 79
Habitat selection:
  Canis latrans, 973
  Canis lupus, 576
  Lynx rufus, 973
  Mephitis mephitis, 1068
  microhabitat affinities of small mammals, 948
Habitat structure:
  Balantiopteryx plicata, 38
  Glaucomys sabrinus griseifrons, 663
Habitat use:
  Cavia aperea, 788
  Glaucomys sabrinus griseifrons, 663
  Glirulus japonicus, 30
  Octodon degus, 260
  Oryzomys palustris, 569
  Tamiasciurus hudsonicus, 225
Hapalemur griseus, 421
Hares:
  Lepus (3 spp.), 591
Hearing:
  Ctenomys sociabilis, 797
  Ctenomys talarum, 290
Heart rate:
  Capreolus capreolus, 245
Heather vole, 97
Heterogeneity bias, 618
Hibernation:
  body surface temperature of Ursus americanus, 414
  Corynorhinus (2 spp.), 989
  Marmota flaviventris, 25
Highveld mole-rat, 1225
Hispid cotton rat, 552, 948, 1077, 1102
Hoarding:
  Callosciurus finlaysonii, 254
  gerbils, 67
Hoary bat, 15, 995
Hodomys alleni, 395
Holographic neural networks, 273
Home range:
  Canis lupus, 576
  Dipodomys heermanni arenae, 1002
  Glirulus japonicus, 30
  Lynx rufus, 983
  Marmota monax, 1087
  Martes pennanti, 649
  Microtus oregoni, 87
  Mustela frenata, 79
  Peromyscus maniculatus, 87
  Procyon lotor, 483
  Sciurus griseus, 511
  Sigmodon hispidus, 1102
  Spermophilus mohavensis, 517
  Thallomys nigricauda, 440
Horses, 611
Huggins closed captures model:
  Ursus arctos, 618
Humpback whale, 963
Hunting strategies/techniques:
  Lampronycteris brachyotis, 1116
  Taxidea taxus, 1019
Hunting success:
  Panthera pardus, 756
Hybrid zone, 1201
Hydrogen isotope, 995
Hydrurga leptonyx, 1215

I
Idaho:
  Taxidea taxus, 1060
Ideal-free distribution, 216, 1053
Illinois:
  Microtus (2 spp.), 216
  Procyon lotor, 483
Immigration:
  Microtus ochrogaster, 781, 1009
India:
  Cynopterus sphinx, 1
  Pteropus giganteus, 1
  Rousettus leschenaulti, 1, 806
Indiana:
  Eptesicus fuscus, 15
  Lasionycteris noctivagans, 15
  Lasiurus (2 spp.), 15
  Myotis (3 spp.), 15
  Nycticeius humeralis, 15
  Pipistrellus subflavus, 15
Indiana myotis, 15, 460
Indonesia:
  Spilocuscus (8 spp.), 825
Infant behavior:
  Microtus agrestis, 282
Infanticide:
  Microtus ochrogaster, 781, 1124
Insectivory:
  Eptesicus fuscus, 460
  Lasiurus borealis, 460
  Myotis (3 spp.), 460
  Nycticeius humeralis, 460
  Pipistrellus subflavus, 460
Interbirth interval:
  Giraffa camelopardalis, 428
Interval censoring, 932
Introduced species:
  Callosciurus finlaysonii, 254
  Crocidura dsinezumi, 396
Isoodon macrourus, 296
Israel:
  Gerbillus (2 spp.), 67
Italy:
  Callosciurus finlaysonii, 254
Ixodes scapularis, 1015

J
Japan:
  Glirulus japonicus, 30
  Ursus thibetanus japonicus, 311
Japanese black bear, 311
Japanese dormouse, 30

K
Kangaroo, 1053
Kangaroo rat:
  Dipodomys heermanni arenae, 1002
  Dipodomys stephensi, 51
Kansas:
  Microtus ochrogaster, 552
  Peromyscus (2 spp.), 552
  Peromyscus maniculatus, 562
  Reithrodontomys megalotis, 552
  Sigmodon hispidus, 552, 1102
  Synaptomys cooperi, 552
Keen's mouse, 540, 1149
Kriging, 1068

L
Lactation:
  Acomys cahirinus, 1106
  Cervus elaphus nelsoni, 714
Lagostomus maximus, 384
Lama guanicoe, 72
Lampronycteris brachyotis, 1116
Landscape:
  badger mounds, 1060
  bat habitat fragmentation, 688
  fragmentation by agriculture, 79
  predator movement in prairie landscape, 187
  ticks and mice in forest fragments, 1015
Land tenure:
  Lynx rufus, 983
Langidium:
  peruanum, 384
  viscacia, 384
  wolffsohni, 384
Larderhoarding, 67
Lasionycteris noctivagans, 15
Lasiurus:
  borealis, 15, 460
  cinereus, 15, 995
Latitude:
  cotton rat life histories and, 1077
  size cline, 331
Latrine behavior:
  lemurs, 420

Least chipmunk, 97
Lemmings, 379, 552
Lemmus trimucronatus, 379
Lemurs, 420
Leopard, 302, 756
Leopard seal, 1215
Lepilemur sp., 421
Leptonychotes weddellii, 1215
Leptonycteris nivalis, 453
Lepus:
  arcticus, 591
  othus, 591
  timidus, 591
Leschenault's rousette, 806
Life history:
  Sigmodon hispidus, 1077
  timing of events, 932
Liomys:
  irroratus, 658
  pictus, 658
Lionycteris, 404
Litter size:
  Microtus socialis guentheri, 748
  Peromyscus leucopus, 940
  Sigmodon hispidus, 1077
Little brown bat (myotis), 15, 273, 460
Livetrapping. *See also* Capture-recapture studies
  Ctenomys minutus, 1201
  Glaucomys sabrinus griseifrons, 663
  Peromyscus maniculatus, 562
  small mammals for richness assessment, 524
Lobodon carinophaga, 1215
Lochophylla:
  chocoana, 404
  handleyi, 404
  mordax, 404
  robusta, 404
  thomasi, 404
Locomotion:
  Isoodon macrourus, 296
Logistic regression, 663, 1068
Long-eared bat, 989
Long-tailed shrew, 331
Long-tailed weasel, 79
Long-term dynamics:
  grassland rodent community, 552
Lyme disease, 1015
Lynx rufus, 973, 983

M
Macrohabitat:
  Glaucomys sabrinus griseifrons, 663
Macropus giganteus, 1053
Madagascar:
  Hapalemur griseus, 421
  Lepilemur sp., 421
Maine:
  Marmota monax, 1087
  Peromyscus leucopus, 743
  Sorex (2 spp.), 331
Major histocompatibility complex (MHC), of Antarctic pinnipeds, 1215
Male genital system:
  Chironectes minimus, 834
Male reproductive success:
  Odocoileus virginianus, 356
Managed forests, 87
Management:
  sexual segregation, implications of, 1039
Mandibular morphology:
  Balaenoptera acutorostrata, 446
  Peromyscus polionotus subgriseus, 1130
Mark-recapture studies:
  Microtus ochrogaster, 1009
  minimum number alive (MNA) population indices, 959
  Sigmodon hispidus, 1102
  Ursus arctos, 618
Marmota:
  flaviventris, 25
  monax, 1087
Marsh rice rat, 569
Marsupials:
  Chironectes minimus, 834
  Isoodon macrourus, 296
  Macropus giganteus, 1053
  Spilocuscus (8 spp.), 825
Martens, 470
Martes:
  americana, 470
  pennanti, 470, 640, 649
Masked shrew, 97, 375
Maternal age:
  Peromyscus leucopus, 940
Maternal behavior:
  Microtus socialis guentheri, 748
  Neotomodon alstoni, 268
Maternal investment:
  Giraffa camelopardalis, 428
Mating:
  Cervus elaphus hispanicus, 761
  Microtus ochrogaster, 1124
  Odocoileus virginianus, 356
  Tadarida braziliensis, 113
  Thallomys nigricauda, 440
Meadow vole, 216
Megadontomys thomasi, 395
Megaptera novaeangliae, 963
Mephitis mephitis, 187, 1068
Meriones tamariscinus, 105
Mesophylla macconnelli, 126
Mesopredator:
  Mephitis mephitis, 1068
Metapopulation:
  Canis lycaon, 625
  Martes pennanti, 640
  Puma concolor, 1207
  Ursus americanus, 140
Mexican long-nosed bat, 453
Mexican prairie dog, 1095
Mexico:
  Baiomys (2 spp.), 162
  Balantiopteryx plicata, 38
  cave use by Mexican bats, 675
  Corynorhinus (2 spp.), 989
  Cratogeomys gymnurus species group, 1170
  Cynomys mexicanus, 1095
  Leptonycteris nivalis, 453
  mammal distribution prediction, 658
  Myotis vivesi, 133
  Neotomodon alstoni, 268
  Peromyscus (6 spp.), 1184
  Peromyscus aztecus, 196
  Reithrodontomys fulvescens, 196
  Ursus americanus, 140
Microclimate, at bat roosts, 675
Microhabitat partitioning, 531
Microhabitat selection:
  Lampronycteris brachyotis, 1116
Microhabitat use:
  affinities of small mammals, 948
  Glaucomys sabrinus griseifrons, 663
Microsatellite DNA:
  Bison bison, 924
  Ctenomys minutus, 1201
  Equus caballus, 611
  Eumetopias jubatus, 338
  Martes pennanti, 640
  Odocoileus virginianus, 356, 768
  Puma concolor, 1207
Microtus:
  agrestis, 282
  montanus, 97
  ochrogaster, 216, 552, 781, 948, 1009, 1124
  oregoni, 87
  pennylvanicus, 216
  socialis guentheri, 748
Middle ear morphology:
  Ctenomys sociabilis, 797
  Ctenomys talarum, 290
Migration:
  Lasiurus cinereus, 995
  Leptonycteris nivalis, 453
  Odocoileus virginianus, 505
Mines, as hibernation roosts, 989
Minimum convex polygon, 511, 517
Minimum number alive (MNA) population indices, 959
Mink, 432
Minke whale, 446
Minnesota:
  Odocoileus virginianus, 505
Mitochondrial DNA studies:
  Artibeus (10 spp.), 866
  Canis lycaon, 625
  chinchilids, 384
  Chiroptera, 321
  Cratogeomys gymnurus species group, 1170
  Crocidura (8 spp.), 396
  Eumetopias jubatus, 338
  Lepus (3 spp.), 591
  Lonchophyllini, 404
  Myotis (2 spp.), 842
  Myotis vivesi, 133
  neotomine-peromycine rodents, 389
  Peromyscus (6 spp.), 1184
  Peromyscus keeni, 1149
  Peromyscus truei species group, 1160
  Scapteromys (2 spp.), 897
  Sorex (2 spp.), 331, 875
  Thrichomys (4 spp.), 316
  Ursus americanus, 140
Mohave ground squirrel, 51
Mole, 120
Molecular divergence:
  chinchilids, 384
  Thrichomys (4 spp.), 316
Molecular evolution:
  Baiomys (2 spp.), 162
Molecular systematics:
  Baiomys (2 spp.), 162
  Myotis vivesi, 133
  neotomine-peromyscine rodents, 1194
  Peromyscus truei species group, 1160
Mole-rat, 1225
Molossidae:
  Eumops (2 spp.), 852
Molt, seasonal:
  Lasiurus cinereus, 995
Mongoose, banded, 491
Monitor lizard, 723
Monogamy:
  Microtus ochrogaster, 1124
Montane shrew, 97, 886
Montane vole, 97
Morphological variation:
  Oxymycterus (3 spp.), 148
  Panthera pardus, 302
  Phocoenoides dalli, 347
  Sorex (3 spp.), 886
  Ursus thibetanus japonicus, 311
Morphology:
  Balaenoptera acutorostrata, 446

Chironectes minimus, 834
Cratogeomys gymnurus species group, 1170
Ctenomys sociabilis, 797
Ctenomys talarum middle ear, 290
Rousettus leschenaulti, 806
Sorex (2 spp.), 331
Morphometrics:
Artibeus (10 spp.), 866
Cryptomys hottentotus pretoriae, 1225
Eumops floridanus, 852
Myotis (2 spp.), 842
Oxymycterus (3 spp.), 148
Panthera pardus, 302
Peromyscus polionotus subgriseus, 1130
Sorex (3 spp.), 886
Mortality:
Sigmodon hispidus, 1077
Spermophilus richardsonii, 1019
Motherhood:
hunting success in Panthera pardus, 756
Mouse:
Aztec, 196
deer, 87, 97, 552, 562, 948
fulvous harvest, 196
Keen's, 540, 1149
oldfield, 1130
spiny, 1106
volcano, 268
western harvest, 552
western jumping, 97
white-footed, 552, 743, 940, 948
Movements:
Cervus elaphus, 774
Isoodon macrourus, 296
Mustela frenata, 79
Odocoileus virginianus, 505
Peromyscus maniculatus, 562
of predators in prairie landscape, 187
Procyon lotor, 483
Sigmodon hispidus, 1102
Spermophilus mohavensis, 517
of voles across habitat boundaries, 216
Mule deer, 498
Multiple paternity:
Mustela vison, 432
Odocoileus virginianus, 356
Mungos mungo, 491
Muscle activity:
hibernating Ursus americanus, 414
Mustela:
frenata, 79
vison, 432
Myotis:
californicus, 842
cilolabrum, 842
lucifugus, 15, 273, 460
septentrionalis, 15, 273, 460
sodalis, 15, 460
vivesi, 133

N
Namibia:
Acinonyx jubatus jubatus, 19
Nasua nasua, 478
Natal dispersal:
Tamiasciurus hudsonicus, 225
Natal philopatry:
Meriones tamariscinus, 105
Natural selection, in Antarctic pinnipeds, 1215
Nectar foraging, by pteropodid bats, 1
Neotoma:
cinerea, 395
floridana, 948
leucodon, 395
mexicana, 395
Neotomine-peromyscine rodents, 389, 1194
Neotomodon alstoni, 268, 395
Nevada:
Equus caballus, 611
Niche theory:
Odocoileus hemionus, 498
sexual segregation and, 1039
North America:
Lasiurus cinereus, 995
Martes pennanti, 640
Myotis (2 spp.), 842
Peromyscus truei species group, 1160
Sorex (2 spp.), 331
Sorex (3 spp.), 886
Taxidea taxus, 633
Thomomys talpoides, 731
Ursus arctos horribilis, 731
North American red squirrel, 225
North Dakota:
Mephitis mephitis, 187
Vulpes vulpes, 187
Northern brown bandicoot, 296
Northern flying squirrel, 206, 663
Northern long-eared bat, 273
Northern myotis, 15, 460
North Pacific:
Megaptera novaeangliae, 963
Northwest Territories:
Canis lupus, 576
Norway:
Castor fiber, 1138
Nuclear DNA analysis:
Cratogeomys (6 spp.), 1170
neotomine-peromyscine rodents, 1194
Nycticeius humeralis, 15, 460
Nyctimene albiventer, 8
Nyctomys sumichrasti, 395

O
Ochrotomys nuttalli, 395
Octodon degus, 260, 384
Odocoileus:
hemionus, 498
virginianus, 356, 505, 658, 768
Ohio:
Microtus ochrogaster, 781, 1009
Oklahoma:
microhabitat affinities of small mammals, 948
Oldfield mouse, 1130
Olfactory communication:
latrine behavior in lemurs, 420
Ommatophoca rossii, 1215
Oncifelis geoffroyi, 1111
Ontogeny, scent gland, 1138
Onychomys:
arenicola, 395
leucogaster, 395
Operational taxonomic unit (OTU):
Dasymys (4 spp.), 911
Orange-bellied bat, 1116
Oregon:
Microtus oregoni, 87
Peromyscus maniculatus, 87
Ornate shrew, 886
Oryzomys:
couesi, 658
palustris, 569
Osgoodomys banderanus, 395
Ototylomys phyllotis, 395
Ovis dalli, 932
Oxymycterus:
dasytrichus, 148
judex, 148
rufus, 148

P
Paleomammalogy:
Dicrostonyx groenlandicus, 379
Eumops floridanus, 852
Lemmus trimucronatus, 379
Panthera pardus, 302
Panama:
Lampronycteris brachyotis, 1116
Panmixia, in cougar populations, 1207
Panthera pardus, 302, 756
Papua New Guinea:
Dobsonia minor, 8
Nyctimene albiventer, 8
Paranyctimene raptor, 8
Pteropus conspicillatus, 8
Spilocuscus (8 spp.), 825
Syconycteris australis, 8
Paraguay:
Artibeus (2 spp.), 688
bat community ecology, 698
bat habitat fragmentation, 688
Carollia perspicillata, 688
Chrotopterus auritus, 688
Glossophaga sorcina, 688
Platyrrhinus lineatus, 688
Pygoderma bilabiatum, 688
Sturnira lilium, 688
Paranyctimene raptor, 8
Parental behavior. *See also* Maternal behavior; Paternal behavior
Microtus socialis guentheri, 748
Parity:
Peromyscus leucopus, 940
Patch selection, 1053
Paternal behavior:
Microtus ochrogaster, 1124
Neotomodon alstoni, 268
Paternity assignment:
Odocoileus virginianus, 356
Pedigree:
Cynomys ludovicianus, 58
Peromyscus:
attwateri, 395, 948, 1160
aztecus, 196
beatae, 395, 1184
boydlii, 1184
californicus, 395
difficilis, 1160
eremicus, 395
gratus, 1160
keeni, 1149
keeni macrorhinus, 540
leucopus, 395, 552, 743, 940, 948, 1015
levipes, 658, 1184
maniculatus, 87, 97, 395, 552, 562, 948
melanophrys, 658
mexicanus, 658
nasutus, 1160
pectoralis, 1160
polionotus subgriseus, 1130
sagax, 1184
schmidlyi, 1184
simulus, 1184
truei, 1160
winkelmanni, 395
zarhynchus, 395
Phalanger:
orientalis, 825
ornatus, 825
Phalangeridae:
Spilocuscus (8 spp.), 825
Phenacomys intermedius, 97
Phocoenoides dalli, 347
Photo identification:
Megaptera novaeangliae, 963

Phyllostomidae:
Lampronycteris brachyotis, 1116
Phylogenetics
arctic organisms, 591
Artibeus (10 spp.), 866
chinchilids, 384
Chiroptera, 321
Crocidura (8 spp.), 396
Myotis (2 spp.), 842
neotomine-peromyscine rodents, 389, 1194
Peromyscus (6 spp.), 1184
Peromyscus truei species group, 1160
Phocoenoides dalli, 347
Sorex (2 spp.), 331, 875
Phylogeny:
Lepus (3 spp.), 591
Lonchophyllini, 404
Myotis vivesi, 133
Panthera pardus, 302
Phocoenoides dalli, 347
Thrichomys (4 spp.), 316
Vampyressa, 126
Phylogeography:
arctic organisms, 591
Eumetopias jubatus, 338
Myotis (2 spp.), 842
Peromyscus keeni, 1149
Scapteromys (2 spp.), 897
Sorex (2 spp.), 875
Sorex (3 spp.), 886
Spermophilus parryii, 601
Ursus americanus, 140
Pinnipeds, 1215
Pipistrellus subflavus, 15, 460
Piscivory:
Myotis vivesi, 133
Pizonyx vivesi, 133
Platalina, 404
Platyrrhinus lineatus, 688
Pliestocene:
lemmings, 379
Pocket gopher, 731, 1170
Pollination, by pteropodid bats, 1
Population density:
Clethrionomys gapperi, 97
Dipodomys stephensi, 51
Glaucomys sabrinus griseifrons, 663
Marmota monax, 1087
Peromyscus leucopus, 743
Peromyscus maniculatus, 97
Phenacomys intermedius, 97
Tamias minimus, 97
Population differentiation:
Ctenomys minutus, 1201
Phocoenoides dalli, 347
Population dynamics:
Glaucomys sabrinus, 206
grassland rodent community, 552
Leptonycteris nivalis, 453
Mungos mungo, 491
Oryzomys palustris, 569
Puma concolor, 1207
Tamiasciurus douglasii, 206
tropical cloud forest bats, 237
Population fragmentation:
Taxidea taxus, 633
Population genetics:
Equus caballus, 611
Population size:
Cynomys ludovicianus, 58
Dipodomys stephensi, 51
minimum number alive (MNA) population indices, 959
Ursus arctos, 618
Population structure:
Martes pennanti, 640
Sorex (2 spp.), 875
Ursus thibetanus japonicus, 311
Prairie, predator movement in, 187
Prairie dogs:
black-tailed, 58
Mexican, 1095
Prairie vole, 216, 552, 781, 1009, 1124
Predation:
effect of refuse feeding, 491
on Rhombomys opimus, 723
risk for Cavia aperea, 788
Predators:
Canis latrans, 973
Genetta genetta, 524
Lynx rufus, 973
Martes (2 spp.), 470
Mephitis mephitis, 187
movements in prairie landscape, 187
Oncifelis geoffroyi, 1111
Panthera pardus, 756
Puma concolor, 1207
Taxidea taxus, 1019
Tyto alba, 524
Vulpes vulpes, 187
Pregnancy:
Acomys cahirinus, 1106
Prey:
of Martes (2 spp.), 470
Rhombomys opimus, 723
Spermophilus richardsonii, 1019
Prey selection:
by insectivorous bats, 460
Primates:
Hapalemur griseus, 421
Lepilemur sp., 421
Prince of Wales flying squirrels, 663
Principal components analysis:
Artibeus species taxonomy, 866
Cryptomys hottentotus pretoriae, 1225
Florida bonneted bat, 852
microhabitat affinities of small mammals, 948
phylogenetics of small-footed myotis, 842
Procyon lotor, 483
Program MARK, 963
Pteropus:
conspicillatus, 8
giganteus, 1
Puma concolor, 1207
Pygoderma bilabiatum, 688

Q
Quebec:
Sorex (2 spp.), 331

R
Raccoon, 483
Radiotelemetry:
Canis latrans, 973
Canis lupus, 576
Capreolus capreolus, 245
Cavia aperea, 788
Cervus elaphus, 774
Dipodomys heermanni arenae, 1002
Glirulus japonicus, 30
Lampronycteris brachyotis, 1116
Lynx rufus, 973
Martes pennanti, 649
Mephitis mephitis, 187
Mustela frenata, 79
Odocoileus virginianus, 505
Procyon lotor, 483
Sciurus griseus, 511
Spermophilus mohavensis, 517
Tamiasciurus hudsonicus, 225
Thallomys nigricauda, 440
Ursus americanus, 414
Ursus arctos, 618
Vulpes vulpes, 187
Rats:
African water, 911
kangaroo, 1002
Oryzomys palustris, 569
Sigmodon hispidus, 552, 948, 1077
Stephens' kangaroo, 51
swamp, 897
Thallomys nigricauda, 440
Recovery, of Cynomys mexicanus, 1095
Red-backed vole, 97
Red bat, 15, 460
Red deer, 761, 774
Red fox, 187
Red squirrel, 225
Red wolf, 625
Refuse feeding, 491
Reithrodontomys:
fulvescens, 196, 395, 948
megalotis, 552, 658
montanus, 948
raviventris, 395
sumichrasti, 395
Relocation:
Martes pennanti, 649
Reproduction:
Acomys cahirinus, 1106
Cervus elaphus hispanicus, 761
Cervus elaphus nelsoni, 714
Cryptomys hottentotus pretoriae, 1225
Microtus ochrogaster, 1124
Microtus socialis guentheri, 748
Mungos mungo, 491
Odocoileus virginianus, 356, 768
Peromyscus leucopus, 940
Spermophilus mohavensis, 517
Reproductive effort/allocation:
Sigmodon hispidus, 1077
Talpa occidentalis, 120
Reproductive strategies:
Giraffa camelopardalis, 428
Mustela vison, 432
Residents:
Lynx rufus, 983
Resource defense:
latrine behavior in lemurs, 420
polygyny in Lama guanicoe, 72
Resource limitation:
Glaucomys sabrinus, 206
Tamiasciurus douglasii, 206
Resource partitioning:
coyotes and bobcats, 973
old world fruit bats, 8
Rest-site use:
Glirulus japonicus, 30
Rhombomys opimus, 723
Richardson's ground squirrel, 1019
Richness:
assessment of small mammal, 524
bat habitat fragmentation, 688
Rodentia:
Abrocoma cinerea, 384
Acomys cahirinus, 1106
Baiomys (2 spp.), 162, 395
Baiomys musculus, 658
Callosciurus finlaysonii, 254
Cavia aperea, 788
Chaetodipus hispidus, 948
Chinchilla (2 spp.), 384
Clethrionomys gapperi, 97
Clethrionomys gapperi wrangeli, 540

Cratogeomys (6 spp.), 1170
Crocidura (8 spp.), 396
Cryptomys hottentotus pretoriae, 1225
Cryptotis magna, 658
Cryptotis parva, 948
Ctenomys minutus, 1201
Ctenomys sociabilis, 797
Ctenomys talarum, 290
Cynomys ludovicianus, 58
Cynomys mexicanus, 1095
Dasymys (4 spp.), 911
Dicrostonyx groenlandicus, 379
Dipodomys heermanni arenae, 1002
Dipodomys stephensi, 51
Gerbillus (2 spp.), 67
Glaucomys sabrinus, 206
Glirulus japonicus, 30
Hodomys alleni, 395
Lagostomus maximus, 384
Langidium (3 spp.), 384
Lemmus trimucronatus, 379
Liomys (2 spp.), 658
Megadontomys thomasi, 395
Meriones tamariscinus, 105
Microtus (2 spp.), 216
Microtus agrestis, 282
Microtus montanus, 97
Microtus ochrogaster, 552, 781, 948, 1009, 1124
Microtus oregoni, 87
Microtus socialis guentheri, 748
Neotoma (3 spp.), 395
Neotoma floridana, 948
neotomine-peromyscine rodents, 389, 1194
Neotomodon alstoni, 268, 395
Nyctomys sumichrasti, 395
Ochrotomys nuttalli, 395
Octodon degus, 260, 384
Onychomys (2 spp.), 395
Oryzomys couesi, 658
Oryzomys palustris, 569
Osgoodomys banderanus, 395
Ototylomys phyllotis, 395
Oxymycterus (3 spp.), 148
Peromyscus (2 spp.), 552
Peromyscus (3 spp.), 658, 948
Peromyscus (6 spp.), 1160, 1184
Peromyscus (8 spp.), 395
Peromyscus aztecus, 196
Peromyscus keeni, 1149
Peromyscus keeni macrorhinus, 540
Peromyscus leucopus, 743, 940
Peromyscus maniculatus, 87, 97, 562
Peromyscus polionotus subgriseus, 1130
Phenacomys intermedius, 97
Reithrodontomys (2 spp.), 948
Reithrodontomys (3 spp.), 395
Reithrodontomys fulvescens, 196
Reithrodontomys megalotis, 552, 658
Rhombomys opimus, 723
Scapteromys (2 spp.), 897
Sciurus aureogaster, 658
Sciurus griseus, 511
Scotinomys xerampelinus, 395
Sigmodon hispidus, 552, 948, 1077, 1102
Sorex (2 spp.), 97, 875
Sorex (3 spp.), 886
Sorex dispar, 331
Sorex gaspensis, 331
Spermophilus (2 spp.), 43
Spermophilus lateralis, 97
Spermophilus mohavensis, 517
Spermophilus parryii, 601
Spermophilus richardsonii, 1019
Synaptomys cooperi, 552
Talpa occidentalis, 120
Tamiasciurus douglasii, 206
Tamiasciurus hudsonicus, 225
Tamias minimus, 97
Thallomys nigricauda, 440
Thomomys talpoides, 731
Thrichomys (4 spp.), 316
Tylomys nudicaudus, 395
Xenomys nelsoni, 395
Zapus princeps, 97
Roe deer, 245
Roost site:
  cave use by Mexican bats, 675
  Corynorhinus (2 spp.), 989
  Leptonycteris nivalis, 453
  Tadarida braziliensis, 113
Ross seal, 1215
Rousettus leschenaulti, 1, 806
Ruminants, sexual segregation in, 1039
Russia:
  Meriones tamariscinus, 105

S
Sac-winged bat, 38
Scapteromys:
  aquaticus, 897
  tumidus, 897
Scatterhoarding, 67
Scent marking:
  Castor fiber, 1138
  Marmota monax, 1087
Sciurus:
  aureogaster, 658
  griseus, 511
Scotinomys xerampelinus, 395
Sea lion, Steller, 338
Seals, 1215
Seasonality:
  Callosciurus finlaysonii, 254
  Capreolus capreolus, 245
  Nasua nasua, 478
  Peromyscus leucopus, 940
  Peromyscus maniculatus, 562
Settlement:
  Tamiasciurus hudsonicus, 225
Sex allocation:
  Odocoileus virginianus, 768
Sex ratio:
  Giraffa camelopardalis, 428
  Odocoileus virginianus, 768
Sexual dimorphism:
  Castor fiber, 1138
Sexual segregation:
  definitions, hypotheses, and implications for conservation and management, 1039
  Odocoileus hemionus, 498
Sexual selection:
  Cervus elaphus hispanicus, 761
Sheppard's correction, 932
Short-tailed field vole, 282
Shrew:
  Crocidura (8 spp.), 396
  Cryptotis magna, 658
  Gaspé, 331
  long-tailed, 331
  masked, 97, 875
  montane, 97, 886
  ornate, 886
  smoky, 875
  wandering, 886
  white-toothed, 396
Shrub-steppe ecosystem, effect of badger mounds, 1060
Sigmodon hispidus, 552, 948, 1077, 1102
Sigmodontinae, 1194
Silver-haired bat, 15
Site fidelity:
  Lama guanicoe, 72
Ski-run development, small mammal survival following, 97
Skull morphology:
  Dasymys (4 spp.), 911
  Panthera pardus, 302
  Peromyscus polionotus subgriseus, 1130
  Ursus thibetanus japonicus, 311
Skunk, striped, 187, 1068
Small-footed myotis, 842
Small mammals:
  microhabitat affinities of small mammals, 948
  microhabitat use, 531
  minimum number alive (MNA) population indices, 959
  richness assessment, 524
  survival and downed wood, 87
  survival following ski-run development, 97
Smoky shrew, 875
Social structure:
  Cavia aperea, 788
  Cryptomys hottentotus pretoriae, 1225
  Cynomys ludovicianus, 58
  Dipodomys heermanni arenae, 1002
  Dipodomys stephensi, 51
  Lynx rufus, 983
  Marmota flaviventris, 25
  Microtus ochrogaster, 781, 1124
  Rhombomys opimus, 723
  Thallomys nigricauda, 440
Social vole, 748
Sonography:
  Spermophilus (2 spp.), 43
Sorex:
  cinereus, 97, 875
  dispar, 331
  fumeus, 875
  gaspensis, 331
  monticolus, 97, 886
  ornatus, 886
  vagrans, 886
South Africa:
  Cryptomys hottentotus pretoriae, 1225
  Panthera pardus, 756
  Thallomys nigricauda, 440
South America. *See also specific countries*
  Abrocoma cinerea, 384
  Chinchilla (2 spp.), 384
  Lagostomus maximus, 384
  Langidium (3 spp.), 384
  Nasua nasua, 478
  Octodon degus, 384
  Oxymycterus (3 spp.), 148
  Scapteromys (2 spp.), 897
South Dakota:
  Cynomys ludovicianus, 58
Southeast Asia:
  Panthera pardus, 302
Southern bog lemming, 552
Southern red-backed vole, 540
Space use:
  Lampronycteris brachyotis, 1116
  Macropus giganteus, 1053
  Odocoileus hemionus, 498
  Sciurus griseus, 511
  Spermophilus mohavensis, 517
  Thallomys nigricauda, 440
Spacing:
  Dipodomys heermanni arenae, 1002
  Marmota monax, 1087
Spain:
  small mammal richness, 524
  Talpa occidentalis, 120

Spatial organization:
Cavia aperea, 788
Mustela vison, 432
Procyon lotor, 483
Spatial scale, 87, 531
Spatial variation:
Dipodomys stephensi, 51
Species boundaries:
Oxymycterus, 148
Sperm:
Odocoileus virginianus, 768
Spermophilus:
lateralis, 43, 97
mohavensis, 517
parryii, 601
richardsonii, 1019
saturatus, 43
Spilocuscus:
kraemeri, 825
maculatus chrysorrhos, 825
maculatus goldiei, 825
maculatus maculatus, 825
maculatus nudicaudatus, 825
papuensis, 825
rufoniger, 825
wilsoni, 825
Spiny mouse, 1106
Spotted cuscus, 825
Squirrels:
Douglas, 206
Finlayson's, 254
Mohave ground, 51
northern flying, 206
Prince of Wales flying, 663
red, 225
Richardson's ground, 1019
Spermophilus parryii, 601
western gray, 511
Stable isotope analysis, 995
Steller sea lion, 338
Stephens' kangaroo rat, 51
Stepping-stone range expansion, 640
Striped skunk, 187, 1068
Sturnira lilium, 688
Subterranean mammals. *See also specific species*
Cryptomys hottentotus pretoriae, 1225
Ctenomys minutus, 1201
Ctenomys sociabilis, 797
Ctenomys talarum, 290
Superfetation and superfecundation:
Mustela vison, 432
Surface temperature:
hibernating Ursus americanus, 414
Survival:
Clethrionomys gapperi, 97
Megaptera novaeangliae, 963
Microtus oregoni, 87
Mungos mungo, 491
Peromyscus maniculatus, 87, 97
Phenacomys intermedius, 97
Rhombomys opimus, 723
Sigmodon hispidus, 1077
Tamias minimus, 97
Swamp rat, 897
Syconycteris australis, 8
Sympatry:
Canis latrans and Lynx rufus, 973
Martes (2 spp.), 470
Oxymycterus, 148
Synaptomys cooperi, 552
Systematics:
Baiomys (2 spp.), 162
Cratogeomys gymnurus species group, 1170
Eumops floridanus, 852
Myotis vivesi, 133
neotomine-peromyscine rodents, 1194
Peromyscus truei species group, 1160
Scapteromys (2 spp.), 897
Spilocuscus (8 spp.), 825
Vampyressa, 126

T
Tadarida braziliensis, 113
Tallgrass prairie, 562
Talpa occidentalis, 120
Tamarisk gerbil, 105
Tamiasciurus:
douglasii, 206
hudsonicus, 225
Tamias minimus, 97
Tampering bias, 959
Taxidea:
taxus, 1019, 1060
taxus jeffersonii, 633
taxus taxus, 633
Taxonomy:
Artibeus (10 spp.), 866
Peromyscus keeni, 1149
Sorex (2 spp.), 331
Spilocuscus (8 spp.), 825
Temperate rainforest:
endemics, 540, 663
flying squirrel microhabitat use, 663
Temperature regulation:
Octodon degus, 260
Temporal variation:
Dipodomys stephensi, 51
Tennessee:
Mephitis mephitis, 1068
Territorial fidelity:
Lama guanicoe, 72
Territoriality:
Castor fiber, 1138
Dipodomys heermanni arenae, 1002
Marmota monax, 1087
Mustela vison, 432
Territory size:
Tamiasciurus hudsonicus, 225
Territory vacancies:
Microtus ochrogaster, 1009
Texas:
Bison bison, 924
Oryzomys palustris, 569
Tadarida braziliensis, 113
Thallomys nigricauda, 440
Thermoregulation, in bats, 675
Thrichomys:
apereoides, 316
inermis, 316
pachyurus, 316
Ticks, 1015
Time budget:
Octodon degus, 260
Time-series analysis, 743
Timing of life-history events, 932
Tool use:
Taxidea taxus, 1019
Tooth wear:
Acinonyx jubatus jubatus, 19
Cryptomys hottentotus pretoriae, 1225
Torpor:
Glirulus japonicus, 30
Trachops cirrhosus, 708
Transients:
Lynx rufus, 983
Travel:
Cervus elaphus, 774
Odocoileus virginianus, 505
Travel cost:
effect on food hoarding, 67
Triphasic growth curves, 120
Tuco-tucos, 290, 797, 1201
Tundra, 576
Tylomys nudicaudus, 395

U
Uganda:
Mungos mungo, 491
Ultrasonic vocalization:
Microtus agrestis, 282
Undermatching, 1053
Unequal intervals, correction for, 932
Ungulates, sexual segregation in, 1039
United States. *See also specific states*
Baiomys (2 spp.), 162
Castor canadensis, 1145
Myotis (2 spp.), 842
Puma concolor, 1207
Sorex (2 spp.), 875
Spermophilus (2 spp.), 43
Taxidea taxus, 1060
Thomomys talpoides, 731
Ursus americanus, 140, 414
Ursus arctos horribilis, 731
Ursus:
americanus, 414
arctos, 618
arctos horribilis, 731
thibetanus japonicus, 311

V
Vampyressa:
bidens, 126
brocki, 126
pusilla, 126
thyone, 126
Vampyriscus:
bidens, 126
brocki, 126
Vampyrum spectrum, 708
Variance, correcting, 932
Vegetation:
flying squirrel microhabitat use, 663
small mammal survival and downed wood, 87
vole movement across habitat boundaries, 216
Virginia:
Sigmodon hispidus, 1077
Vocalization:
alarm call variability in golden-mantled ground squirrels, 43
mating in Tadarida braziliensis, 113
ultrasonic in Microtus agrestis, 282
Volcano mouse, 268
Vole:
creeping, 87
heather, 97
meadow, 216
montane, 97
prairie, 216, 552, 781, 1009, 1121
red-backed, 97
short-tailed field, 282
social, 748
southern red-backed, 540
Vulpes vulpes, 187

W
Wandering shrew, 886
Washington:
Sciurus griseus, 511
Water balance:
Acomys cahirinus, 1106

Water rat, 911
Weasel, long-tailed, 79
Weddell seal, 1215
Western gray squirrel, 511
Western harvest mouse, 552
Western jumping mouse, 97
Wetlands, 569
Whales:
  Balaenoptera acutorostrata, 446
  Megaptera novaeangliae, 963
White-footed mouse, 395, 552, 743, 940, 948, 1015
White-tailed deer, 356, 505, 768
Wing morphology:
  Rousettus leschenaulti, 806
Wolf, 576, 625
Woodchuck, 1087
Wyoming:
  Cervus elaphus nelsoni, 714

X
Xenomys nelsoni, 395

Y
Yapok, 834
Yellow-bellied marmot, 25
Yellowstone:
  Cervus elaphus nelsoni, 714
  Thomomys talpoides, 731
  Ursus arctos horribilis, 731
Yellowstone elk, 714
Yinpterochiroptera, 321

Z
Zapus princeps, 97
Zinc-finger protein DNA regions, 1145